arXiv:1610.00845v1 [cs.IT] 4 Oct 2016

Isometrically Self-dual Cyclic Codes

Yun Fan and Liang Zhang
Dept of Mathematics, Central China Normal University, Wuhan 430079, China

October 5, 2016

Abstract

General isometries of cyclic codes, including multipliers and translations, are introduced; and isometrically self-dual cyclic codes are defined. In terms of Type-I duadic splittings given by multipliers and translations, a necessary and sufficient condition for the existence of isometrically self-dual cyclic codes is obtained. A program to construct isometrically self-dual cyclic codes is provided, and illustrated by several examples. In particular, a class of isometrically self-dual MDS cyclic codes, which are alternant codes from a class of generalized Reed-Solomon codes, is presented.

Keywords: cyclic code, isometry, isometrically self-dual, MDS cyclic codes.

MSC2010: 12E20, 94B60.

1 Introduction

Self-dual codes and cyclic codes are two important and long-time topics which are interesting in both theoretical perspective and technological practice.

Let $\mathbb{F}_q$ be a finite field with q elenments, where q is a power of a prime. Cyclic codes of length n over $\mathbb{F}_q$ can be described as ideals of the quotient algebra $\mathbb{F}_q[X]/\langle X^n-1\rangle$ of the polynomial algebra $\mathbb{F}_q[X]$ with respect to the ideal $\langle X^n-1\rangle$ generated by X^n-1. If the greatest common divisor $\gcd(q,n)=1$, then $\mathbb{F}_q[X]/\langle X^n-1\rangle$ is a semisimple algebra. Semisimple cyclic codes are studied and applied extensively. However, it is a pity that there are no (euclidean) self-dual semisimple cyclic codes. An alternative research is the study on *duadic* cyclic codes, which are initiated by Leon, Masley and Pless [18], and developed extensively from various directions, e.g., see [21, 24, 5, 22, 25, 9, 11, 26, 19, 20, 23, 2, 15, 12, 17, 14].

In the semisimple case, the irreducible factors in $\mathbb{F}_q[X]$ of X^n-1 correspond one-to-one to the *q-cyclotomic cosets* (*q-cosets* in short) of the residue ring $\mathbb{Z}_n$

Email addresses: yfan@mail.ccnu.edu.cn (Y. Fan)

of the integer ring $\mathbb{Z}$ modulo n. By $\mathbb{Z}_n^*$ we denote the subset of the elements $s \in \mathbb{Z}_n$ such that $\gcd(s, n) = 1$. For any $s \in \mathbb{Z}_n^*$, we have a permutation μ_s of $\mathbb{Z}_n$ such that (see [16, p.138]):

$$\mu_s(i) = si \pmod{n}, \qquad \forall\ i \in \mathbb{Z}_n. \tag{1.1}$$

The permutation μ_s is called a *multiplier*. Correspondingly, we have an isomorphism $\hat{\mu}_s$ of the algebra $\mathbb{F}_q[X]/\langle X^n - 1\rangle$ such that (see [16, Eqn (4.4)]):

$$\hat{\mu}_s\big(a(X)\big) = a(X^s) \pmod{X^n - 1}, \qquad \forall\ a(X) \in \mathbb{F}_q[X]/\langle X^n - 1\rangle. \tag{1.2}$$

The isomorphism $\hat{\mu}_s$ preserves the Hamming weight obviously (the isomorphisms which preserve Hamming weight are called *isometries*). If there is a subset $P \subseteq \mathbb{Z}_n$ which is a union of some q-cosets such that $\mathbb{Z}_n = \{0\} \bigcup P \bigcup \mu_s(P)$ is a partition, then P and $\mu_s(P)$ are called an even-like duadic splitting of $\mathbb{Z}_n$. In that case, $\mathbb{F}_q[X]/\langle X^n - 1\rangle = C + \hat{\mu}_s(C)$ and $C \bigcap \hat{\mu}_s(C)$ is the cyclic code generated by the vector $(1, \cdots, 1)$, where C is the cyclic code with defining set P. Please refer to [16, Ch. 6] for details.

Constacyclic codes are a natural generalization of cyclic codes, they are ideals of the quotient algebra $\mathbb{F}_q[X]/\langle X^n - \lambda\rangle$, where $\lambda \in \mathbb{F}_q^*$ is a non-zero element. Semisimple constacyclic codes are still the most important case. In the semisimple case, [10] showed that self-dual negacyclic codes (i.e., $\lambda = -1$) exist. Blackford [3] proved that semisimple self-dual constacyclic codes do not exist except for the negacyclic case; and he showed conditions that the self-dual negacyclic codes exist. A more concise condition for the existence of self-dual negacyclic codes appeared in [7, Corollary 21].

An innovation to develop further the theory is the concept of *iso-dual constacyclic codes* introduced in [4]. Assume that λ is a primitive r-th root of unity (of course, r divides $q - 1$). Similarly to the cyclic codes, the set of roots of $X^n - \lambda$ corresponds to a subset of $\mathbb{Z}_{nr}$, denoted by $1 + r\mathbb{Z}_{nr}$, which consists of the elements $i \in \mathbb{Z}_{nr}$ such that $i \equiv 1 \pmod{r}$. The irreducible factors of $X^n - \lambda$ correspond bijectively to the q-cosets on the set $1 + r\mathbb{Z}_{nr}$. For $s \in \mathbb{Z}_{nr}^*$, Blackford [4] defined a multiplier μ_s on $1 + r\mathbb{Z}_{nr}$ similarly to Eqn (1.1), and defined an isometry φ_s of $\mathbb{F}_q[X]/\langle X^n - \lambda\rangle$ similarly to Eqn (1.2). Then, a constacyclic code C is said to be *iso-dual* if there is an isometry which maps C onto its dual code $C^\perp$. If there is a subset $P \subseteq (1 + r\mathbb{Z}_{nr})$ which is a union of some q-cosets such that $1 + r\mathbb{Z}_{nr} = P \bigcup \mu_s(P)$ is a partition, then P and $\mu_s(P)$ are called a *Type-I duadic splitting* of $1 + r\mathbb{Z}_{nr}$ given by μ_s; in that case, $\mathbb{F}_q[X]/\langle X^n - 1\rangle = C \oplus \varphi_s(C)$ is a direct sum, where C is the constacyclic code with defining set P. Blackford [4] proved that iso-dual constacyclic codes correspond one to one to the Type-I duadic splittings given by multipliers; and showed conditions for the existence of such splittings (an issue was addressed in [6]). A necessary and sufficient condition for the existence of the Type-I duadic splittings given by multipliers was presented in [7, Corollary 19]. With these notations, however, the iso-dual cyclic codes do not exist either, because any multiplier μ_s on $\mathbb{Z}_n$ leaves 0 invariant, hence the Type-I duadic splittings of $\mathbb{Z}_n$ given by multipliers do not exist; see also Remark 3.3 (ii) below.

It is an interesting question: whether we can find a generalized self-duality which occurs in semisimple cyclic codes? The key idea is that, in essence, an isometry of cyclic codes is an algebra isomorphism preserving Hamming weight; though multipliers are isometries, there might be isometries outside of the multipliers, and those isometries might give isometrically self-dual cyclic codes. We should investigate the general isometries of cyclic codes to discover the isometries besides multipliers, then we would answer the question positively.

In Section 2, we start with an investigation of the general isometries of semisimple cyclic codes of length n, and define the *isometrically self-dual cyclic codes*, abbreviated as *iso-self-dual cyclic codes*. The permutations of $\mathbb{Z}_n$ corresponding the general isometries include not only multipliers, and translations as well. Then, we transform the study of iso-self-dual semisimple cyclic codes into the study of Type-I duadic splittings of $\mathbb{Z}_n$ given by the multipliers and translations.

In Section 3, we present necessary and sufficient conditions for the existence of Type-I duadic splittings of $\mathbb{Z}_n$ given by multipliers and translations. Meanwhile, a method of construction of such splittings is also provided.

In Section 4, we exhibit a program to construct iso-self-dual semisimple cyclic codes, and illustrate it with examples. If the length n is even and $n/2$ is odd, then for any finite field $\mathbb{F}_q$ with q coprime to n the iso-self-dual cyclic codes of length n over $\mathbb{F}_q$ exist and the construction of them is interesting. In particular, a class of iso-self-dual MDS cyclic codes, which are alternant codes from a class of generalized Reed-Solomon codes, is obtained.

2 Isometrically self-dual cyclic codes

Let $\mathbb{F}_q$ be the finite field with cardinality $|\mathbb{F}_q| = q$, where q is a prime power, let $\mathbb{F}_q^*$ be the set of non-zero elements of $\mathbb{F}_q$. Let n be a positive integer.

By $\mathbb{Z}_n$ we denote the residue ring of the integer ring $\mathbb{Z}$ modulo n, and by $\mathbb{Z}_n^*$ we denote the multiplicative group consisting of units of $\mathbb{Z}_n$.

By $R_n = \mathbb{F}_q[X]/\langle X^n - 1\rangle$ we denote the quotient algebra of the polynomial algebra $\mathbb{F}_q[X]$ modulo the ideal $\langle X^n - 1\rangle$ generated by $X^n - 1$. Any element of R_n has a unique representative of degree $< n$: $a(X) = a_0 + a_1X + \cdots + a_{n-1}X^{n-1}$, which is identified with the word $\mathbf{a} = (a_0, a_1, \cdots, a_{n-1}) \in \mathbb{F}_q^n$. By $\mathrm{w}(a(X))$ we denote the Hamming weight of the word $\mathbf{a}$. Any ideal C of R_n is called a *cyclic code* of length n over $\mathbb{F}_q$.

In the following we *always* assume that $\gcd(q, n) = 1$, i.e., the cyclic codes considered in the following are all semisimple. We always assume that θ is a primitive n-th root of unity (in a suitable extension of $\mathbb{F}_q$). Then

$$X^n - 1 = \prod_{i \in \mathbb{Z}_n} (X - \theta^i). \tag{2.1}$$

In this way, roots of $X^n - 1$ correspond bijectively to elements of $\mathbb{Z}_n$.

Definition 2.1. A transformation $\varphi : R_n \to R_n$ is called an *isometry* of R_n if φ is an algebra automorphism of R_n and preserves the Hamming weight of R_n (i.e., $\mathrm{w}(\varphi(a(X))) = \mathrm{w}(a(X))$ for all $a(X) \in R_n$).

General isometries between constacyclic codes were introduced in [8] to classify constacyclic codes, with less attention paid to cyclic codes. We need precise characterizations of isometries of cyclic codes as follows.

Lemma 2.2. *A transformation $\varphi : R_n \to R_n$ is an isometry if and only if there is a positive integer $s \in \mathbb{Z}_n^*$ and an integer $t \in \mathbb{Z}_n$ satisfying that $qt \equiv t \pmod n$ such that*

$$\varphi\big(a(X)\big) = a(\theta^t X^s) \pmod{X^n - 1}, \qquad \forall\ a(X) = \sum_{i=0}^{n-1} a_i X^i \in R_n. \tag{2.2}$$

Proof. Assume that $\varphi : R_n \to R_n$ is an isometry. Since φ preserves the Hamming weight, there is an element $b \in \mathbb{F}_q^*$ and a positive integer $s \in \mathbb{Z}_n$ such that $\varphi(X) = bX^s$. Because φ is an algebra automorphism, the n elements:

$$\varphi(X^i) = (bX^s)^i = b^i X^{si} \pmod{X^n - 1}, \qquad i = 0, 1, \cdots, n-1, \tag{2.3}$$

are a basis of R_n. If the greatest common divisor $d = \gcd(s, n) > 1$, then the remainder of si modulo n is a multiple of d; hence the n elements in Eqn (2.3) are not a basis of R_n, which is not the case. So the integer s is coprime to n, i.e., $s \in \mathbb{Z}_n^*$. Since $X^n = 1$ in R_n, we have

$$1 = \varphi(X^n) = \varphi(X)^n = b^n X^{sn} = b^n;$$

that is, $b = \theta^t$ for an integer $t \in \mathbb{Z}_n$. By the Galois theory, $\theta^t \in \mathbb{F}_q$ if and only if $(\theta^t)^q = \theta^t$ which is equivalent to that $qt \equiv t \pmod n$. Then Eqn (2.2) holds obviously.

Conversely, assume that $s \in \mathbb{Z}_n^*$, $t \in \mathbb{Z}_n$ satisfying that $qt \equiv t \pmod n$, and Eqn (2.2) holds. Note that $\theta^t \in \mathbb{F}_q$ because $qt \equiv t \pmod n$. For any $a(X), b(X) \in R_n$, in R_n we have the following computation:

$$\varphi\big(a(X)b(X)\big) = a(\theta^t X^s)b(\theta^t X^s) = \varphi\big(a(X)\big)\varphi\big(b(X)\big) \pmod{X^n - 1}.$$

Similarly, we can check that $\varphi\big(a(X) + b(X)\big) = \varphi\big(a(X)\big) + \varphi\big(b(X)\big)$. Thus, φ is an algebra endomorphism of R_n. For $i = 0, 1, \cdots, n-1$, there are integers h_i and r_i with $0 \le r_i \le n-1$ such that $si = nh_i + r_i$. Then

$$\varphi(X^i) = \theta^{ti} X^{r_i}, \qquad i = 0, 1, \cdots, n-1.$$

Since s is coprime to n, it is easy to see that $i \mapsto r_i$ is a permutation of $\{0, 1, \cdots, n-1\}$. Let P_s be the corresponding permutation matrix. By D_t we denote the diagonal matrix with $1, \theta^t, \cdots, \theta^{t(n-1)}$ on the diagonal positions. Then $\varphi(X^i) = \theta^{ti} X^{r_i}$ implies that:

- *With respect to the basis* $1, X, \cdots, X^{n-1}$ *of* R_n*, the matrix of* φ *is the monomial matrix* $D_t P_s$*.*

Hence φ is an algebra automorphism and preserves the Hamming weight. □

Remark 2.3. For $s \in \mathbb{Z}_n^*$ and $t \in \mathbb{Z}_n$ with $qt \equiv t \pmod{n}$, we have a positive integer $s^{-1} \in \mathbb{Z}_n^*$ such that $ss^{-1} \equiv 1 \pmod{n}$; and for $-t$ we still have $q(-t) \equiv -t \pmod{n}$. By $\varphi_{s,t}$ we denote the isometry of R_n defined by

$$\varphi_{s,t}: \quad R_n \longrightarrow R_n, \quad a(X) \longrightarrow a(\theta^{-t} X^{s^{-1}}) \pmod{X^n - 1}. \tag{2.4}$$

That is, $\varphi_{s,t}\big(a(X)\big) = a(\theta^{-t} X^{s^{-1}}) \pmod{X^n - 1}$, it is the unique polynomial of degree $< n$ congruent to $a(\theta^{-t} X^{s^{-1}})$ modulo $X^n - 1$. We illustrate two specific cases.

(i) If $s = 1$. then we take $s^{-1} = 1$ and, for any $a(X) = \sum_{i=0}^{n-1} a_i X^i \in R_n$, since $\deg a(\theta^{-t} X) = \deg a(X) < n$, we have

$$\varphi_{1,t}\big(a(X)\big) = a(\theta^{-t} X) = a_0 + \theta^{-t} a_1 X + \cdots + \theta^{-t(n-1)} a_{n-1} X^{n-1}. \tag{2.5}$$

(ii) If $s = -1$ and $t = 0$, then we take $s^{-1} = n - 1$ as $(-1)(n-1) \equiv 1 \pmod{n}$; since $X^n \equiv 1 \pmod{X^n - 1}$, by Eqn (2.4), for $a(X) = \sum_{i=0}^{n-1} a_i X^i$ we have

$$\varphi_{-1,0}\big(a(X)\big) \equiv X^n \sum_{i=0}^{n-1} a_i X^{(n-1)i} \equiv \sum_{i=0}^{n-1} a_i X^{n-i} \pmod{X^n - 1};$$

i.e.,

$$\varphi_{-1,0}\big(a(X)\big) = a_0 + a_{n-1} X + \cdots + a_1 X^{n-1}. \tag{2.6}$$

As usual, by $C^\perp$ we denote the (Euclidean) dual code of any code C. It is easy to check that, for any cyclic code $C \subseteq R_n$, its dual code $C^\perp \subseteq R_n$ is still a cyclic code.

Definition 2.4. For a cyclic code $C \subseteq R_n$ over $\mathbb{F}_q$ of length n, if there is an isometry $\varphi_{s,t}$ of R_n such that $\varphi_{s,t}(C) = C^\perp$, then we say that C is an *isometrically self-dual cyclic code*, or, an *iso-self-dual cyclic code* for short.

Thus, an iso-self-dual cyclic code C has the same algebraic structure as its dual code $C^\perp$; specifically, $h(X)$ is a check polynomial of C if and only if $\varphi_{s,t}\big(h(X)\big)$ is a check polynomial of $\varphi_{s,t}(C) = C^\perp$. Meanwhile, an iso-self-dual cyclic code C has the same weight structure as $C^\perp$; specifically, C and $C^\perp$ have the same weight distributions; in other words, C is a *formally self-dual* cyclic code, see [16, page 378].

In the rest of this section we translate the study of iso-self-dual cyclic codes into a study of corresponding permutations on the index set $\mathbb{Z}_n$.

For any pair s, t where $s \in \mathbb{Z}_n^*$ and $t \in \mathbb{Z}_n$ satisfying that $qt \equiv t \pmod{n}$, it is easy to see that the map

$$\rho_{s,t}: \ \mathbb{Z}_n \longrightarrow \mathbb{Z}_n, \ \ i \longrightarrow s(i+t) \pmod{n}, \tag{2.7}$$

is a permutation of $\mathbb{Z}_n$, which we call a *q-permutation of $\mathbb{Z}_n$*.

We list some related known facts in the following remark.

Remark 2.5. (i) Note that, if $t = 0$, then we denote $\mu_s = \rho_{s,0}$, which is called a *multiplier* on $\mathbb{Z}_n$ in literature. On the other hand, if $s = 1$, then we denote $\tau_t = \rho_{1,t}$. Since $\tau_t(i) = i + t \pmod{n}$ for $i \in \mathbb{Z}_n$ and t should satisfy that "$qt \equiv t \pmod{n}$" where q is involved, we call τ_t a *q-translation* on $\mathbb{Z}_n$. Obviously, $\rho_{s,t} = \mu_s \tau_t$.

(ii) Since $q \in \mathbb{Z}_n^*$, we have a special multiplier μ_q, and the μ_q-orbits (i.e., the orbits on $\mathbb{Z}_n$ of the permutation group $\langle \mu_q \rangle$ generated by μ_q) are usually named *q-cyclotomic cosets*. In the following, we call μ_q-orbits on $\mathbb{Z}_n$ by *q-cosets*, and use $\mathbb{Z}_n/\mu_q$ to stand for the quotient sets consisting of all q-cosets on $\mathbb{Z}_n$.

(iii) For any q-coset $Q \in \mathbb{Z}_n/\mu_q$, the polynomial $f_Q(X) = \prod_{i \in Q}(X - \theta^i)$ is irreducible in $\mathbb{F}_q[X]$. We have the monic irreducible decomposition: $X^n - 1 = \prod_{Q \in \mathbb{Z}_n/\mu_q} f_Q(X)$. Obviously, a subset $P \subseteq \mathbb{Z}_n$ is μ_q-invariant (i.e., $\mu_q(P) = P$) if and only if $P = \bigcup_{j=1}^k Q_j$ is a union of some q-cosets $Q_1, \cdots, Q_k$. If it is the case, then the polynomial $f_P(X) = \prod_{i \in P}(X - \theta^i) = \prod_{j=1}^k f_Q(X)$ is a monic divisor of $X^n - 1$. In this way, the monic divisors of $X^n - 1$ correspond one to one to the μ_q-invariant subsets of $\mathbb{Z}_n$.

(iv) Each μ_q-invariant subset P of $\mathbb{Z}_n$ corresponds to exactly one cyclic code (i.e., ideal) of R_n, denoted by C_P, such that the polynomial $f_P(X)$ is a check polynomial of C_P; both the degree $\deg f_P(X)$ and the dimension $\dim C_P$ equal to the cardinality $|P|$ of P. The correspondence $P \mapsto C_P$ is a lattice isomorphism between the lattice of μ_q-invariant subsets of $\mathbb{Z}_n$ and the lattice of cyclic codes of R_n.

Lemma 2.6. *Let $s \in \mathbb{Z}_n^*$ and $t \in \mathbb{Z}_n$ satisfying that $qt \equiv t \pmod{n}$, and s^{-1} be a positive integer such that $ss^{-1} \equiv 1 \pmod{n}$. Then for $f_Q(X)$, where $Q \in \mathbb{Z}_n/\mu_q$ is a q-coset, we have an $\mathbb{F}_q$-polynomial $u_Q(X)$ which is coprime to $X^n - 1$ and has leading coefficient $\theta^{-t|Q|}$ such that*

$$f_Q(\theta^{-t} X^{s^{-1}}) = u_Q(X) f_{\rho_{s,t}(Q)}(X).$$

Proof. Since $qt \equiv t \pmod{n}$, we see that $\theta^{-t} \in \mathbb{F}_q$ and

$$\rho_{s,t}(Q) = s(Q + t) = \{s(i + t) \pmod{n} \mid i \in Q\}$$

is still a q-coset, and the cardinalities $\big|Q\big| = \big|\rho_{s,t}(Q)\big|$. Then:

$$\begin{aligned} f_Q(\theta^{-t} X^{s^{-1}}) &= \prod_{i \in Q}(\theta^{-t} X^{s^{-1}} - \theta^i) = \prod_{i \in Q} \theta^{-t}\big(X^{s^{-1}} - (\theta^{(i+t)s})^{s^{-1}}\big) \\ &= \theta^{-t|Q|} \prod_{i \in Q} \frac{X^{s^{-1}} - (\theta^{(i+t)s})^{s^{-1}}}{X - \theta^{(i+t)s}} \cdot \prod_{i \in Q}(X - \theta^{(i+t)s}). \end{aligned}$$

Since the map $Q \to \rho_{s,t}(Q)$, $i \mapsto s(i+t) \pmod n$, is a bijection, we see that

$$\prod_{i\in Q}(X-\theta^{(i+t)s}) = \prod_{i\in \rho_{s,t}(Q)}(X-\theta^{i}) = f_{\rho_{s,t}(Q)}(X).$$

It remains to show that $\prod_{i\in Q}\frac{X^{s^{-1}}-(\theta^{(i+t)s})^{s^{-1}}}{X-\theta^{(i+t)s}}$ is coprime to X^n-1. To do it, it is enough to show that any root θ^j of X^n-1 is not a root of the polynomial $\frac{X^{s^{-1}}-(\theta^{(i+t)s})^{s^{-1}}}{X-\theta^{(i+t)s}}$ for any $i \in Q$. If $j\equiv(i+t)s \pmod n$, then $\theta^j = \theta^{(i+t)s}$ which is not a root of the polynomial $\frac{X^{s^{-1}}-(\theta^{(i+t)s})^{s^{-1}}}{X-\theta^{(i+t)s}}$. Otherwise, $j\not\equiv(i+t)s \pmod n$ hence $\theta^j \neq \theta^{(i+t)s}$; since s^{-1} is coprime to n, $(\theta^j)^{s^{-1}} - (\theta^{(i+t)s})^{s^{-1}} \neq 0$; i.e., θ^j is not a root of the polynomial $\frac{X^{s^{-1}}-(\theta^{(i+t)s})^{s^{-1}}}{X-\theta^{(i+t)s}}$. □

Corollary 2.7. *Let s, t, s^{-1} be as in Lemma 2.6, and $P \subseteq \mathbb{Z}_n$ be a μ_q-invariant subset. Then we have an $\mathbb{F}_q$-polynomial $u_P(X)$ which is coprime to X^n-1 and has leading coefficient $\theta^{-t|P|}$ such that*

$$f_P(\theta^{-t}X^{s^{-1}}) = u_P(X) f_{\rho_{s,t}(P)}(X).$$

Proof. There are some q-cosets $Q_1, \cdots, Q_k$ such that P is a disjoint union $P = \bigcup_{i=1}^k Q_i$. Then $\rho_{s,t}(P)$ is a disjoint union $\rho_{s,t}(P) = \bigcup_{i=1}^k \rho_{s,t}(Q_i)$, and $f_P(X) = \prod_{i=1}^k f_{Q_i}(X)$. By Lemma 2.6, $f_{Q_i}(\theta^{-t}X^{s^{-1}}) = u_{Q_i}(X) f_{\rho_{s,t}(Q_i)}(X)$ with $u_{Q_i}(X)$ being coprime to X^n-1 and having leading coefficient $\theta^{-t|Q_i|}$. So

$$f_P(\theta^{-t}X^{s^{-1}}) = \prod_{i=1}^k u_{Q_i}(X) \cdot \prod_{i=1}^k f_{\rho_{s,t}(Q_i)}(X) = \prod_{i=1}^k u_{Q_i}(X) \cdot f_{\rho_{s,t}(P)}(X),$$

where $\prod_{i=1}^k u_{Q_i}(X)$ is coprime to X^n-1 and has leading coefficient $\theta^{-t|P|}$. □

Though the polynomials $\varphi_{s,t}\big(f_P(X)\big)$ and $f_{\rho_{s,t}(P)}(X)$ are different from each other in general, by the above result, we can find the polynomial $f_{\rho_{s,t}(P)}(X)$ from the polynomial $\varphi_{s,t}\big(f_P(X)\big)$.

Theorem 2.8. *Let $s \in \mathbb{Z}_n^*$ and $t \in \mathbb{Z}_n$ satisfying that $qt \equiv t \pmod n$, let P be a μ_q-invariant subset of $\mathbb{Z}_n$. Then*

$$f_{\rho_{s,t}(P)}(X) = \gcd\big(f_P(\theta^{-t}X^{s^{-1}}),\, X^n-1\big) = \gcd\big(\varphi_{s.t}\big(f_P(X)\big),\, X^n-1\big).$$

Proof. From Corollary 2.7, we get immediately that

$$f_{\rho_{s,t}(P)}(X) = \gcd\big(f_P(\theta^{-t}X^{s^{-1}}),\, X^n-1\big).$$

Note that $\varphi_{s,t}\big(f_P(X)\big)$ is the remainder of $f_P(\theta^{-t}X^{s^{-1}})$ modulo X^n-1. The greatest common divisor in the right hand side of the above equality is clearly equal to $\gcd\big(\varphi_{s.t}\big(f_P(X)\big),\, X^n-1\big)$. □

We describe two specific cases of the theorem. Recall that, for a polynomial $a(X) = a_0 + a_1X + \cdots + a_dX^d$, the polynomial $X^d \cdot a(X^{-1}) = a_d + a_{d-1}X + \cdots + a_0X^d$ is called the *reciprocal polynomial* of $a(X)$.

Corollary 2.9. *(i)* $f_{\rho_{1,t}(P)}(X) = \theta^{t|P|} f_P(\theta^{-t}X)$.

(ii) Let $a_0 = (-1)^{|P|}\prod_{i\in P}\theta^i$ *be the constant term of* $f_P(X)$. *Then*

$$f_{\rho_{-1,0}(P)}(X) = f_{-P}(X) = a_0^{-1} \cdot X^{|P|} f_P(X^{-1}).$$

Proof. Note that both $f_P(X)$ and $f_{\rho_{s,t}(P)}(X)$ are monic divisors of X^n-1 with degree $|P|$.

(i). In this case where $s = 1$ (hence $s^{-1} = 1$), the polynomial $u_P(X)$ in Corollary 2.7 is in fact the constant $\theta^{-t|P|}$.

(ii). Note that $\rho_{-1,0}(P) = (-1)(P + 0) = -P = \{-i \mid i \in P\}$. Denote $|P| = d$. We can write $f_P(X) = a_0 + a_1X + \cdots + a_dX^d$, where $a_d = 1$ and $a_0 = (-1)^d\prod_{i\in P}\theta^i$ because $f_P(X) = \prod_{i\in P}(X - \theta^i)$. By Eqn (2.6),

$$\begin{aligned}\varphi_{-1,0}\big(f_P(X)\big) &= a_0 + a_1X^{n-1} + \cdots + a_dX^{n-d}\\ &= X^{n-d}(a_0X^d + a_1X^{d-1} + \cdots + a_d) - a_0(X^n - 1)\\ &= X^{n-d} \cdot X^d f_P(X^{-1}) - a_0(X^n - 1).\end{aligned}$$

Since θ^i for $i \in (-P)$ are all roots of $X^d f_P(X^{-1})$ (cf. [16, Lemma 4.4.7]), $X^d f_P(X^{-1})$ is a divisor of $X^n - 1$. Thus

$$\gcd\big(\varphi_{-1.0}\big(f_P(X)\big),\, X^n - 1\big) = a_0^{-1}(a_0X^d + a_1X^{d-1} + \cdots + a_d).$$

By Theorem 2.8, we obtain the conclusion (ii). □

We turn to cyclic codes and state the following concise result.

Theorem 2.10. *For any q-invariant subset* $P \subseteq \mathbb{Z}_n$ *and any isometry* $\varphi_{s,t}$

$$\varphi_{s,t}(C_P) = C_{\rho_{s,t}(P)}.$$

Proof. Since $f_P(X)$ is a check polynomial of C_P and $\varphi_{s,t}$ is an algebra automorphism of R_n, the cyclic code $\varphi_{s,t}(C_P)$ has a check polynomial $\varphi_{s,t}\big(f_P(X)\big) = f_P(\theta^{-t}X^{s^{-1}})$ (mod X^n-1). By Corollary 2.7, $u_P(X)f_{\rho_{s,t}(P)}(X)$ is a check polynomial of $\varphi_{s,t}(C_P)$, where $u_P(X)$ is a unit of R_n. So, $f_{\rho_{s,t}(P)}(X)$ is also a check polynomial of $\varphi_{s,t}(C_P)$. By definition (see Remark 2.5 (iv)), $\varphi_{s,t}(C_P) = C_{\rho_{s,t}(P)}$. □

Similar to [4, Theorem 4] and [13, Lemma 3.5], we have the following result.

Lemma 2.11. *Let* P *be a q-invariant subset of* $\mathbb{Z}_n$. *Then*

$$C_P^{\perp} = \varphi_{-1,0}(C_{\bar{P}}) = C_{-\bar{P}}\,,$$

where $\bar{P}{=}\mathbb{Z}_n\backslash P$ *is the complement of* P *in* $\mathbb{Z}_n$.

Proof. By Theorem 2.10, $\varphi_{-1,0}(C_{\bar{P}}) = C_{\rho_{-1,0}(\bar{P})} = C_{-\bar{P}}$.

Let $a(X) = \sum_{i=0}^{n-1} a_i X^i \in C_{\bar{P}}$. For any $c(X) = \sum_{i=0}^{n-1} c_i X^i \in C_P$, set $b(X) = X^{n-1}c(X) \pmod{X^n - 1}$. Then $b(X) \in C_P$ (as C_P is an ideal) and

$$b(X) = c_1 + c_2 X + \cdots + c_{n-1}X^{n-2} + c_0 X^{n-1}.$$

Since $f_P(X)f_{\bar{P}}(X) = X^n - 1$, $a(X)b(X) \equiv 0 \pmod{X^n - 1}$. Observing the coefficient of X^{n-1} in $a(X)b(X) \pmod{X^n - 1}$, we obtain that

$$a_0c_0 + a_1c_{n-1} + \cdots + a_{n-2}c_2 + a_{n-1}c_1 = 0.$$

By Eqn (2.6),

$$\varphi_{-1,0}\big(a(X)\big) = a_0 + a_{n-1}X + \cdots + a_1 X^{n-1}.$$

So the (Euclidean) inner product

$$\big\langle \varphi_{-1,0}\big(a(X)\big), c(X) \big\rangle = a_0c_0 + a_{n-1}c_1 + \cdots + a_1c_{n-1} = 0$$

Thus $\varphi_{-1,0}\big(a(X)\big) \in C_{\bar{P}}^{\perp}$. In conclusion, $\varphi_{-1,0}(C_{\bar{P}}) \subseteq C_{\bar{P}}^{\perp}$. Finally,

$$\dim \varphi_{-1,0}(C_{\bar{P}}) = \dim C_{\bar{P}} = |\bar{P}| = n - |P| = n - \dim C_P = \dim C_{\bar{P}}^{\perp};$$

from which we further get that $\varphi_{-1,0}(C_{\bar{P}}) = C_{\bar{P}}^{\perp}$. □

Theorem 2.12. *Let P be a q-invariant subset of $\mathbb{Z}_n$. The cyclic code $C_P \subseteq R_n$ is an iso-self-dual cyclic code if and only if there is a positive integer $s \in \mathbb{Z}_n^*$ and an integer $t \in \mathbb{Z}_n$ satisfying that $qt \equiv t \pmod{n}$ such that $\mathbb{Z}_n = P \bigcup \rho_{s,t}(P)$ is a partition. If it is the case, then the following hold:*

(i) $\varphi_{-s,t}(C_P) = C_{\bar{P}}^{\perp}$.

(ii) *$h(X)$ is a check polynomial of C_P if and only if $\varphi_{-s,t}(h(X))$ is a check polynomial of $C_{\bar{P}}^{\perp}$ (equivalently, $\varphi_{-s,t}(h(X))$ is a generator polynomial of C_P).*

(iii) *C_P and $C_{\bar{P}}^{\perp}$ have the same weight distributions.*

Proof. Let $\varphi_{s,t}$ be an isometry of R_n. By Theorem 2.10 and Lemma 2.11, $\varphi_{s,t}(C_P) = C_{\bar{P}}^{\perp}$ is equivalent to $C_{\rho_{s,t}(P)} = C_{-\bar{P}}$, i.e., $-\bar{P} = \rho_{s,t}(P) = s(P + t)$. Thus $\varphi_{s,t}(C_P) = C_{\bar{P}}^{\perp}$ if and only if $\bar{P} = -s(P + t) = \rho_{-s,t}(P)$. The latter one is equivalent to that $\mathbb{Z}_n = P \bigcup \rho_{-s,t}(P)$ is a partition of $\mathbb{Z}_n$.

Replacing $-s$ by s in the above argument, we obtain the conclusions (i). Since $\varphi_{-s,t}$ is an algebra automorphism of R_n and preserves Hamming weight, (ii) and (iii) are obtained immediately. □

Definition 2.13. (i) Let $s \in \mathbb{Z}_n^*$, $t \in \mathbb{Z}_n$ satisfying that $qt \equiv t \pmod{n}$, and P be a μ_q-invariant subset of $\mathbb{Z}_n$. If $\mathbb{Z} = P \bigcup \rho_{s,t}(P)$ is a partition of $\mathbb{Z}_n$, then we say that $P, \rho_{s,t}(P)$ are a *Type-I duadic splitting of* $\mathbb{Z}_n$ given by $\rho_{s,t}$.

(ii) If there are s, t, P such that the above (i) holds, then we say that Type-I duadic splittings of $\mathbb{Z}_n$ exist.

Thus, the study of iso-self-dual cyclic codes is translated to the study of Type-I duadic splittings of $\mathbb{Z}_n$. Theorem 2.12 says that iso-self-dual cyclic codes of length n exist if and only if Type-I duadic splittings of $\mathbb{Z}_n$ exist; and, in that case, the information of the iso-self-dual cyclic codes can be obtained from the information of the Type-I duadic splittings of $\mathbb{Z}_n$.

3 Type-I duadic splittings of $\mathbb{Z}_n$

Keep the notations in Section 2. For convenience, we denote

$$G_{q,n} = \big\{\rho_{s,t} \,\big|\, s \in \mathbb{Z}_n^*,\ t \in \mathbb{Z}_n \text{ satisfying that } qt \equiv t \pmod n\big\}, \tag{3.1}$$

where $\rho_{s,t}$ is the q-permutation of $\mathbb{Z}_n$ defined in Eqn (2.7). It is easy to check that

$$\rho_{s',t'}\rho_{s,t} = \rho_{s's,t+s^{-1}t'}, \qquad \forall\ \rho_{s',t'}, \rho_{s,t} \in G_{q,n};$$

in particular,

$$\rho_{s,t}^{-1} = \rho_{s^{-1},-st}, \qquad \forall\ \rho_{s,t} \in G_{q,n}.$$

So, $G_{q,n}$ is a permutation group of $\mathbb{Z}_n$, which we call the *q-permutation group* of $\mathbb{Z}_n$. Recall from Remark 2.5 (i) that we denote $\mu_s = \rho_{s,0}$ and call it a *multiplier*; we denote $\tau_t = \rho_{1,t}$ and call it a *q-translation* on $\mathbb{Z}_n$; and $\rho_{s,t} = \mu_s\tau_t$.

From the condition $qt \equiv t \pmod n$, we have $\tau_t\mu_q = \mu_q\tau_t$. So the subgroup $\langle\mu_q\rangle$ generated by μ_q is a central subgroup of $G_{q,n}$. Then the group $G_{q,n}$ acts on the quotient set $\mathbb{Z}_n/\mu_q$ consisting of q-cosets. The properties of this action determine the existence and constructions of the Type-I duadic splittings of $\mathbb{Z}_n$. For fundamentals of group actions, please refer to [1]. Some elementary results on group actions will be cited later which we list in the following remark; cf. [1] or [7, Lemma 6, Lemma 7] for details.

Remark 3.1. Let a finite group G act on a finite set X. For $\rho \in G$, by $\langle\rho\rangle$ we denote the subgroup generated by ρ. Any orbit of the group $\langle\rho\rangle$ on X is abbreviated as a ρ-orbit.

(i) Let $\rho \in G$. Then there is a subset $S \subseteq X$ such that $X = S \bigcup \rho(S)$ is a partition of X if and only if the length of any ρ-orbit on X is even. If it is the case, for each ρ-orbit $(x_1, x_2, \cdots, x_{2m})$, i.e., $\rho(x_i) = x_{i+1}$ for $1 \le i < 2m$ and $\rho(x_{2m}) = x_1$, we can place $x_1, x_3, \cdots, x_{2m-1}$ into S; once this is done for every ρ-orbit, we obtain a subset $S \subseteq X$ such that $X = S \bigcup \rho(S)$ is a partition.

(ii) The length of any ρ-orbit on X is a divisor of the order of ρ in the group G. In particular, if the order of ρ is a power of 2, then there is a subset $S \subseteq X$ such that $X = S \bigcup \rho(S)$ is a partition of X if and only if there is no element of X which is fixed by ρ.

(iii) Let another finite group G' act on a finite set X'. Then the finite group $G \times G'$ acts on the finite set $X \times X'$. For $(\rho, \rho') \in G \times G'$ and $(x.x') \in X \times X'$, the length of the (ρ, ρ')-orbit on $X \times X'$ containing (x, x') equals to the least common multiple of the lengths of the ρ-orbit on X containing x and the length of the ρ'-orbit on X' containing x'.

Lemma 3.2. *Let $\rho_{s,t} \in G_{q,n}$. A q-coset $Q \in \mathbb{Z}_n/\mu_q$ is fixed by $\rho_{s,t}$ if and only if there is a $k \in Q$ and an integer $j \geq 0$ such that $(q^j - s)k \equiv st \pmod{n}$.*

Proof. Note that, since $\rho_{s,t}(Q)$ is still a q-coset, $\rho_{s,t}(Q) = Q$ if and only if $\rho_{s,t}(Q) \bigcap Q \neq \emptyset$. Thus, $\rho_{s,t}(Q) = Q$ if and only if there is a $k \in Q$ and an integer $j \geq 0$ such that $\rho_{s,t}(k) = q^j k$. The latter is just $(q^j - s)k \equiv st \pmod{n}$. □

Remark 3.3. (i) By $\nu_2(m)$ we denote the 2-adic valuation of any non-zero integer m, i.e., $2^{\nu_2(m)}$ is the largest power of 2 which divides m. For convenience, we appoint that $\nu_2(0) = \infty$.

(ii) If $t = 0$, then "$k = 0$" is always a solution for $(q^j - s)k \equiv st \pmod{n}$; hence the Type-I duadic splittings of $\mathbb{Z}_n$ given by any multiplier $\rho_{s,0} = \mu_s$ do not exist. Thus, in the following we consider the case where $t \not\equiv 0 \pmod{n}$.

(iii) From Lemma 3.4 till to Theorem 3.8, we consider the case where $n = 2^v$ with $v \geq 1$. In that case, q must be odd since we have assumed that $\gcd(q, n) = 1$, and, with the notation in (i), the condition that $qt \equiv t \pmod{n}$ in Eqn (3.1) (i.e. $(q-1)t \equiv 0 \pmod{2^v}$) can be rewritten as $v \leq \nu_2(q-1) + \nu_2(t)$.

Lemma 3.4. *Let $n = 2^v$ with $v \geq 1$. Let $\rho_{s,t} \in G_{q,2^v}$ with $t \not\equiv 0 \pmod{2^v}$. The Type-I duadic splittings of $\mathbb{Z}_{2^v}$ given by $\rho_{s,t}$ exist if and only if*

$$\nu_2(q^j - s) > \nu_2(t), \qquad \forall\ j \geq 0. \tag{3.2}$$

Proof. Since two integers in a residue class of $\mathbb{Z}_{2^v}$ have different 2-adic valuations, we need to show that the inequality (3.2) is independent of the choices of the integers s, t from their residue classes modulo 2^v. Suppose that $s \in \mathbb{Z}_{2^v}^*$ satisfies the inequality (3.2), i.e., $\nu_2(q^j - s) > \nu_2(t)$. For any $s' \equiv s \pmod{2^v}$, we write $s' = s + 2^v k$. Since $0 \leq \nu_2(t) < v$, we get

$$\nu_2(q^j - s') = \nu_2(q^j - s - 2^v k) \geq \min\{\nu_2(q^j - s), v\} > \nu_2(t).$$

So, s' still satisfies the inequality (3.2). Similarly, if $\nu_2(q^j - s) > \nu_2(t)$ and $t' = t + 2^v k$, then $\nu_2(t') = \nu_2(t)$ since $\nu_2(t) < v$, hence $\nu_2(q^j - s) > \nu_2(t')$.

It is easy to see that the group $G_{q,2^v}$ is of order power of 2, hence the order of $\rho_{s,t}$ is a power of 2. Thus the Type-I duadic splittings of $\mathbb{Z}_{2^v}$ given by $\rho_{s,t}$ exist if and only if there is no q-coset of $\mathbb{Z}_{2^v}$ which is fixed by $\rho_{s,t}$, please see Remark 3.1 (ii).

Suppose that the condition (3.2) holds. Then for any $j \geq 0$,

$$\nu_2((q^j - s)k) \geq \nu_2(q^j - s) > \nu_2(t) = \nu_2(st), \qquad \forall\ k \in \mathbb{Z}_n.$$

Since $t \not\equiv 0 \pmod{2^v}$, for any $k \in \mathbb{Z}_n$ we get that $(q^j - s)k \not\equiv st \pmod{2^v}$. By Lemma 3.2, there is no q-coset $Q \in \mathbb{Z}_n/\mu_q$ which is fixed by $\rho_{s,t}$.

Conversely, assume that there is a $j \geq 0$ such that $\nu_2(q^j - s) \leq \nu_2(t) = \nu_2(st)$. Then we can find an integer $k \in \mathbb{Z}_{2^v}$ such that $(q^j - s)k \equiv st \pmod{2^v}$. By Lemma 3.2, the q-coset Q_k containing k is fixed by $\rho_{s,t}$. □

Corollary 3.5. *Let $n = 2^v$ with $v \geq 1$. Let $\rho_{1,t} = \tau_t \in G_{q,2^v}$ with $\nu_2(t) < v$ be a q-translation. The Type-I duadic splittings of $\mathbb{Z}_{2^v}$ given by τ_t exist if and only if*

$$\nu_2(t) < \nu_2(q-1).$$

Proof. If the Type-I duadic splittings of $\mathbb{Z}_{2^v}$ given by τ_t exist, by Lemma 3.4, $\nu_2(t) < \nu_2(q^j - 1)$ for any $j \geq 0$. Taking $j = 1$, we obtain $\nu_2(t) < \nu_2(q-1)$.

Conversely, assume that $\nu_2(t) < \nu_2(q-1)$. Since $(q-1) \mid (q^j - 1)$ for any $j \geq 0$, we get $\nu_2(t) < \nu_2(q^j - 1)$ for any $j \geq 0$. By Lemma 3.4, the Type-I duadic splittings of $\mathbb{Z}_{2^v}$ given by τ_t exist. □

Lemma 3.6. *Let $n = 2^v$ with $v \geq 1$, and $s \in \mathbb{Z}_{2^v}^*$. There is a $t \in \mathbb{Z}_{2^v}$ with $qt \equiv t \pmod{2^v}$ such that Type-I duadic splittings of $\mathbb{Z}_{2^v}$ given by $\rho_{s,t}$ exist if and only if*

$$\nu_2(q^j - s) + \nu_2(q-1) > v, \qquad \forall\ j \geq 0. \tag{3.3}$$

Proof. First we need to show that the inequality (3.3) is independent of the choice of the integer s from its residue class modulo 2^v. Suppose that $s \in \mathbb{Z}_{2^v}^*$ satisfies the inequality (3.3), i.e., $\nu_2(q^j - s) > v - \nu_2(q-1)$. For any $s' \equiv s \pmod{2^v}$ we can write $s' = s + 2^v k$. Since $\nu_2(q-1) > 0$ (recall that q is odd), we get

$$\nu_2(q^j - s') = \nu_2(q^j - s - 2^v k) \geq \min\{\nu_2(q^j - s), v\} > v - \nu_2(q-1).$$

That is, s' still satisfies the inequality (3.3).

If $\nu_2(q-1) \geq v$, by the condition "$\nu_2(q-1)+\nu_2(t) \geq v$" (see Remark 3.3(iii)), we can take a $t \in \mathbb{Z}_{2^v}$ such that $\nu_2(t) = 0$, i.e., t is odd. Since both q and s are odd, for any $j \geq 0$, $\nu_2(q^j - s) \geq 1 > \nu_2(t)$; hence the condition (3.3) is satisfied, and by Lemma 3.4, Type-I duadic splittings of $\mathbb{Z}_{2^v}$ given by $\rho_{s,t}$ exist. In a word, the conclusion of the lemma holds in this case.

Next, we assume that $\nu_2(q-1) < v$, i.e., $0 < v - \nu_2(q-1) < v$.

Suppose that the condition (3.3) is satisfied. Since $\nu_2(q-1) < v$, we can take a $t \in \mathbb{Z}_{2^v}$ such that $\nu_2(t) = v - \nu_2(q-1)$, see Remark 3.3(iii). Then

$$\nu_2(q^j - s) > v - \nu_2(q-1) = \nu_2(t), \qquad \forall\ j \geq 0;$$

by Lemma 3.4, the Type-I duadic splittings of $\mathbb{Z}_{2^v}$ given by $\rho_{s,t}$ exist.

Suppose that the condition (3.3) is not satisfied, i.e., there is a $j \geq 0$ such that $\nu_2(q^j - s) \leq v - \nu_2(q-1)$. For any $\rho_{s,t} \in G_{q,n}$, by Remark 3.3(iii), $\nu_2(q-1) + \nu_2(t) \geq v$. So, $v - \nu_2(q-1) \leq \nu_2(t) < v$. Then $\nu_2(q^j - s) \leq v - \nu_2(q-1) \leq \nu_2(t)$. By Lemma 3.4, Type-I duadic splittings of $\mathbb{Z}_{2^v}$ given by $\rho_{s,t}$ do not exist. □

Lemma 3.7. *Let $n = 2^v$ with $v \geq 1$, let $\rho_{s,t} \in G_{q,n}$. If the Type-I duadic splittings of $\mathbb{Z}_{2^v}$ given by $\rho_{s,t}$ exist, then the Type-I duadic splittings of $\mathbb{Z}_{2^v}$ given by $\rho_{1,t} = \tau_t$ exist.*

Proof. By Lemma 3.4, for $\rho_{s,t}$ the condition (3.2) is satisfied. If $s \equiv 1 \pmod{2^v}$, then we are done. In the following we assume that $s \not\equiv 1 \pmod{2^v}$. For $j = 0$, from the condition (3.2) we get

$$\nu_2(1-s) > \nu_2(t).$$

Taking $j = 1$ we further get $\nu_2(q-s) > \nu_2(t)$, which we can rewrite as follows:

$$\nu_2\big((q-1)+(1-s)\big) > \nu_2(t).$$

If $\nu_2(q-1) < \nu_2(1-s)$, then

$$\nu_2(q-1) = \nu_2\big((q-1)+(1-s)\big) > \nu_2(t).$$

Otherwise, $\nu_2(q-1) \geq \nu_2(1-s)$; since $\nu_2(1-s) > \nu_2(t)$, we still get

$$\nu_2(q-1) > \nu_2(t).$$

For any $j \geq 0$, noting that $(q-1) \mid (q^j-1)$, we obtain that

$$\nu_2(q^j-1) \geq \nu_2(q-1) > \nu_2(t).$$

By Lemma 3.4, the Type-I duadic splittings of $\mathbb{Z}_{2^v}$ given by $\rho_{1,t}$ exist. □

Theorem 3.8. *Let $n = 2^v$. Then the following three statements are equivalent to each other:*

(i) *The Type-I duadic splittings of $\mathbb{Z}_{2^v}$ exist.*

(ii) *The Type-I duadic splittings of $\mathbb{Z}_{2^v}$ given by q-translations exist.*

(iii) $0 < v < 2 \cdot \nu_2(q-1)$.

If it is the case, then there is an integer u such that

$$\max\{0,\ v-\nu_2(q-1)\} \leq u < \min\{v,\ \nu_2(q-1)\}; \tag{3.4}$$

hence $\rho_{1,2^u} = \tau_{2^u} \in G_{q,2^v}$ is a q-translation and the Type-I duadic splittings of $\mathbb{Z}_{2^v}$ given by τ_{2^u} exist.

Proof. Obviously, any one of the three statements implies that $v \geq 1$.

(i) $\Leftrightarrow$ (ii). This follows from Lemma 3.7 immediately.

(ii) $\Rightarrow$ (iii). By Lemma 3.6, (ii) implies that

$$\nu_2(q^j-1) + \nu_2(q-1) > v, \qquad \forall\ j \geq 0.$$

Taking $j = 1$, we get $\nu_2(q-1) + \nu_2(q-1) > v$; i.e., (iii) holds.

(iii) $\Rightarrow$ (ii). Assume that (iii) holds, i.e., $0 < v$ and $v - \nu_2(q-1) < \nu_2(q-1)$. Since q is odd, $0 < \nu_2(q-1)$ and $v - \nu_2(q-1) < v$. Hence

$$0 \leq \max\{0,\ v-\nu_2(q-1)\} < \min\{v,\ \nu_2(q-1)\}.$$

Thus the integer u satisfying Eqn (3.4) exists. Set $t = 2^u$, then $\nu_2(t) = u$. Because $v - \nu_2(q-1) \le u = \nu_2(t)$, by Remark 3.3 (iii), $\tau_t = \rho_{1,t} \in G_{q,2^v}$ is a q-translation. Since Eqn (3.4) implies that $\nu_2(t) < \nu_2(q-1)$, by Corollary 3.5, Type-I duadic splittings of $\mathbb{Z}_{2^v}$ given by τ_t exist. □

The following lemma is a bridge from the case where $n = 2^v$ to the general case where n is any positive integer coprime to q.

Lemma 3.9. *Let $v = \nu_2(n)$ and $n = 2^v n'$ with $2 \nmid n'$, let $\rho_{s,t} \in G_{q,n}$. The following two statements are equivalent to each other:*

(i) The Type-I duadic splittings of $\mathbb{Z}_n$ given by $\rho_{s,t}$ exist.

(ii) The Type-I duadic splittings of $\mathbb{Z}_{2^v}$ given by $\rho_{s,t}$ exist.

Proof. By Chinese Remainder Theorem, we have a natural isomorphism

$$\mathbb{Z}_n \overset{\text{CRT}}{=\!=} \mathbb{Z}_{n'} \times \mathbb{Z}_{2^v} \tag{3.5}$$

which maps any $k \in \mathbb{Z}_n$ to $\big(k \pmod{n'},\, k \pmod{2^v}\big) \in \mathbb{Z}_{n'} \times \mathbb{Z}_{2^v}$. Obviously, $s \in \mathbb{Z}_{n'}^*$ and $s \in \mathbb{Z}_{2^v}^*$; $qt \equiv t \pmod{n'}$ and $qt \equiv t \pmod{2^v}$. That is, we also have the q-permutation $\rho_{s,t}$ of $\mathbb{Z}_{n'}$ and of $\mathbb{Z}_{2^v}$. Since the cardinality of $\mathbb{Z}_{n'}$ is $|\mathbb{Z}_{n'}| = n'$ which is odd, at least one of the $\rho_{s,t}$-orbits on $\mathbb{Z}_{n'}/\mu_q$ has odd length. By Remark 3.1 (iii), the length of any $\rho_{s,t}$-orbit on $\mathbb{Z}_n/\mu_q$ is even if and only if the length of any $\rho_{s,t}$-orbit on $\mathbb{Z}_{2^v}/\mu_q$ is even. By Remark 3.1 (i), the statements (i) and (ii) are equivalent to each other. □

By Lemma 3.9, all the previous results on $\mathbb{Z}_{2^v}$ also hold for the general case $\mathbb{Z}_n$. In particular, from Theorem 3.8 we can get the following result.

Theorem 3.10. *Let $n = 2^{\nu_2(n)} n'$ (hence $2 \nmid n'$) be any positive integer coprime to q. Then the following three statements are equivalent to each other:*

(i) *The Type-I duadic splittings of $\mathbb{Z}_n$ exist.*

(ii) *The Type-I duadic splittings of $\mathbb{Z}_n$ given by q-translations exist.*

(iii) $0 < \nu_2(n) < 2 \cdot \nu_2(q-1)$.

If it is the case, then there is an integer u such that

$$\max\{0,\ \nu_2(n) - \nu_2(q-1)\} \le u < \min\{\nu_2(n),\ \nu_2(q-1)\}; \tag{3.6}$$

hence $\rho_{1,2^u n'} = \tau_{2^u n'} \in G_{q,n}$ is a q-translation and the Type-I duadic splittings of $\mathbb{Z}_n$ given by $\tau_{2^u n'}$ exist.

Proof. Let $v = \nu_2(n)$, i.e., $n = 2^v n'$ with $2 \nmid n'$.

(i) ⇒ (iii). By (i) we assume that $\rho_{s,t} \in G_{q,n}$ such that the Type-I duadic splittings of $\mathbb{Z}_n$ given by $\rho_{s,t}$ exist. By Lemma 3.9, the Type-I duadic splittings of $\mathbb{Z}_{2^v}$ given by $\rho_{s,t}$ exist. By Theorem 3.8, $0 < v < 2 \cdot \nu_2(q-1)$; i.e., (iii) holds.

(iii) $\Rightarrow$ (ii). By Theorem 3.8, (iii) implies that there is an integer u which satisfies Eqn (3.6). It is obvious that $q \cdot 2^u n' \equiv 2^u n' \pmod{n'}$. By Eqn (3.6), $v - \nu_2(q-1) \le u = \nu_2(2^u n')$, i.e., $q \cdot 2^u n' \equiv 2^u n' \pmod{2^v}$, cf. Remark 3.3 (iii). So $q \cdot 2^u n' \equiv 2^u n' \pmod{2^v n'}$ as 2^v and n' are coprime. Recalling that $n = 2^v n'$, we get a q-translation $\rho_{1,2^v n'} = \tau_{2^u n'} \in G_{q,n}$ of $\mathbb{Z}_n$. By Eqn (3.6) again, $\nu_2(2^u n') = u < \nu_2(q-1)$. By Corollary 3.5, the Type-I duadic splitiings of $\mathbb{Z}_{2^v}$ given by $\tau_{2^u n'}$ exist. By Lemma 3.9, the Type-I duadic splitiings of $\mathbb{Z}_n$ given by $\rho_{1,2^v n'} = \tau_{2^u n'}$ exist.

(ii) $\Rightarrow$ (i). Trivial. □

4 Constructions of iso-self-dual cyclic codes

4.1 A program of construction of iso-self-dual cyclic codes

From Theorem 3.10 and Theorem 2.12, we get immediately a necessary and sufficient condition for the existence of iso-self-dual cyclic codes.

Theorem 4.1. *Iso-self-dual cyclic codes over $\mathbb{F}_q$ of length n exist if and only if $0 < \nu_2(n) < 2 \cdot \nu_2(q-1)$.*

Remark 4.2. More precisely, Theorem 3.10, Theorem 2.12 and their arguments provide a program to construct iso-self-dual cyclic codes over $\mathbb{F}_q$ of length n.

S1. (determination of existence of iso-self-dual cyclic codes)

if $\nu_2(n) = 0$ or $\nu_2(n) \ge 2 \cdot \nu_2(q-1)$, then output "`iso-self-dual cyclic codes over` $\mathbb{F}_q$ `of length` n `do not exist`";

otherwise take an integer u satisfying that

$$\max\{0,\ \nu_2(n) - \nu_2(q-1)\} \le u < \min\{\nu_2(n),\ \nu_2(q-1)\},$$

set $n' = n/2^{\nu_2(n)}$, $t = 2^u n'$ and go to S2.

S2. (construction of iso-self-dual cyclic codes)

1) compute q-cosets on $\mathbb{Z}_n$ to get the quotient set $\mathbb{Z}_n/\mu_q$;
2) compute the τ_t-orbits on $\mathbb{Z}_n/\mu_q$;
3) for every τ_t-orbit $(Q_1, Q_2, \cdots, Q_{2m})$ on $\mathbb{Z}_n/\mu_q$ (the length of any τ_t-orbit must be even), place $Q_1, Q_3, \cdots, Q_{2m-1}$ into the subset P (cf. Remark 3.1 (i));
4) output the cyclic code $C_P \subseteq R_n$ with check polynomial $f_P(X)$ and the dual code $C_P^\perp = \varphi_{-1,t}(C_P) = C_{-\bar{P}}$.

Example 4.3. Let $q = 3$, $n = 8$. Then $\nu_2(n) = 3 > 2 = 2\nu_2(q-1)$; so there are no iso-self-dual cyclic codes over $\mathbb{F}_3$ of length 8. In fact, in this case,

$$\mathbb{Z}_8/\mu_3 = \big\{\{0\},\ \{4\},\ \{1,3\},\ \{2,6\},\ \{5,7\}\big\}$$

which consists of 5 q-cosets; any permutation on $\mathbb{Z}_8/\mu_3$ has at least one orbit of odd length.

Example 4.4. Let $q = 5$, $n = 8$. Then $\nu_2(n) = 3 < 4 = 2\nu_2(q-1)$; the condition in Theorem 4.1 is satisfied. We can take $u = 1$:

$$\max\{0,\ \nu_2(n) - \nu_2(q-1)\} = 1 \le 1 < 2 = \min\{\nu_2(n),\ \nu_2(q-1)\}.$$

Then $t = 2$. The set of q-cosets

$$\mathbb{Z}_8/\mu_5 = \big\{\{0\},\ \{4\},\ \{2\},\ \{6\},\ \{1,5\},\ \{3,7\}\big\}.$$

The orbits of the q-translation $\tau_t = \tau_2$ are as follows:

$$\big(\{0\},\ \{2\},\ \{4\},\ \{6\}\big), \quad \big(\{1,5\},\ \{3,7\}\big).$$

We put $\{0\},\ \{4\},\ \{1,5\}$ into P, and get $P = \{0,1,4,5\}$. Then $\rho_{1,t}(P) = \bar{P}$ is the complement of P in $\mathbb{Z}_n$. Take θ to be a primitive 8-th root of unity such that $\theta^2 = 2$. Then C_P is an iso-self-dual cyclic code over $\mathbb{F}_5$ of length 8 with the check polynomial $f_P(X)$ as follows (note that $\theta^5 = -\theta$):

$$\begin{aligned} f_P(X) &= (X-\theta^0)(X-\theta^4)(X-\theta)(X-\theta^5) \\ &= (X-1)(X+1)(X^2-2) \\ &= X^4 + 2X^2 + 2. \end{aligned}$$

And $C_P^{\perp} = \varphi_{-1,t}(C_P)$. Thus $f_{\rho_{-1,t}(P)}(X)$ is a generator polynomial of C_P. By Corollary 2.9, $f_{\rho_{-1,t}(P)}(X)$ can be obtained from $f_P(X)$ as follows (note that $\theta^{-t} = 2^{-1} = 3$):

$$f_{\rho_{1,t}(P)}(X) = \theta^{t|P|} f_P(\theta^{-t}X) = X^4 - 2X^2 + 2;$$

and

$$f_{\rho_{-1,t}(P)}(X) = f_{\rho_{-1,0}(\rho_{1,t}(P))}(X) = 2^{-1}(1 - 2X^2 + 2X^4) = X^4 - X^2 - 2.$$

4.2 The case where $\nu_2(n) = 1$

For the length n, there is a specific case where the iso-self-dual cyclic codes of length n always exist.

Theorem 4.5. *Assume that n is an even positive integer such that $n' = n/2$ is odd. Then, for any odd prime power q coprime to n, the length of any $\tau_{n'}$-orbit on the quotient set $\mathbb{Z}_n/\mu_q$ of q-cosets equals to 2; in particular, Type I duadic splittings of $\mathbb{Z}_n$ given by $\tau_{n'}$ exist.*

Proof. For any $i \in \mathbb{Z}_n$, $\tau_{n'}^2(i) = i + 2n' \equiv i \pmod{n}$. That is, the order of the permutation $\tau_{n'}$ equals to 2; Then the length of any $\tau_{n'}$-orbit on $\mathbb{Z}_n/\mu_q$ equals to either 2 or 1. Suppose a $\tau_{n'}$-orbit on $\mathbb{Z}_n/\mu_q$ has length 1; i.e., consists of only one q-coset Q; then $\tau_{n'}(Q) = Q$ and by Lemma 3.2 there is a $k \in Q$ and an integer $j \ge 0$ such that $(q^j - 1)k \equiv n' \pmod{n}$; this is impossible because both $q^j - 1$ and n are even but n' is odd. Therefore, any $\tau_{n'}$-orbit on $\mathbb{Z}_n/\mu_q$ has length 2. By Remark 4.2, taking any one q-coset from each $\tau_{n'}$-orbit on $\mathbb{Z}_n/\mu_q$ to put into P, we get a partition $\mathbb{Z}_n = P \bigcup \tau_{n'}(P)$. □

Remark 4.6. Let notations be as in Theorem 4.5. In that case we can omit the step S1 in Remark 4.2; moreover, we have a corresponding polynomial version of the step S2 in Remark 4.2. To state it, we first describe two operations on polynomials, which correspond to the q-permutations $\tau_{n'}$ and μ_{-1} respectively.

Let $a(X) = X^d + a_{d-1}X^{d-1} + \cdots + a_0$ be a monic polynomial of degree $\deg a(X) = d$ with non-zero constant a_0.

- $\widetilde{a}(X) = (-1)^{\deg a(X)}a(-X) = X^d - a_{d-1}X^{d-1} + \cdots + (-1)^d a_0$, called the *alternating polynomial* of $a(X)$;
- $a^*(X) = a_0^{-1}X^{\deg a(X)}a(X^{-1})$, called the *monic reciprocal polynomial* of $a(X)$;

For a primitive n-th root θ of unity, $\theta^{-n'} = -1$. Then, by Corollary 2.9, the program S2 in Remark 4.2 can be translated as follows:

S2. (construction of iso-self-dual cyclic codes)

1) decompose $X^n - 1$ into a product of monic irreducible factors;
2) partition the monic irreducible factors of $X^n - 1$ into pairs such that in every pair one polynomial is alternating of the other one;
3) for every pair of the partition in 2), select one factor from the pair;
4) set $h(X)$ to be the product of the factors which are selected in 3); output the cyclic code $C \subseteq R_n$ with check polynomial $h(X)$, and the dual code $C^\perp$ with check polynomial $\widetilde{h}^*(X)$.

We illustrate the above construction with an example.

Example 4.7. Let $n = 10$, $q = 3$. By the above theorem the Type-I duadic splittings of $\mathbb{Z}_{10}$ given by the 3-translation τ_5 exist. By the program in Remark 4.2, we can get one of them as follows. The set of q-cosets

$$\mathbb{Z}_{10}/\mu_3 = \big\{\{0\},\ \{5\},\ \{1,3,9,7\},\ \{2,6,8,4\}\big\}. \tag{4.1}$$

The τ_5-orbits on $\mathbb{Z}_{10}/\mu_3$ are:

$$\big(\{0\},\ \{5\}\big), \quad \big(\{1,3,9,7\},\ \{2,6,8,4\}\big). \tag{4.2}$$

For every τ_5-orbit, select one q-coset as follows:

$$\{0\}, \quad \{1,3,9,7\}. \tag{4.3}$$

Place them into the subset P to get

$$P = \{0,\ 1,\ 3,\ 9 = -1,\ 7 = -3\}, \tag{4.4}$$

and $\tau_5(P) = \{2, 4, 5, 6, 8\} = \bar{P}$,

$$\mu_{-1}\tau_5(P) = -\bar{P} = \{-2, -4, -5, -6, -8\} = \bar{P}. \tag{4.5}$$

Corresponding to the equations (4.1)–(4.5), we list the polynomial versions described in Remark 4.6 as follows.

The monic irreducible decomposition

$$X^n - 1 = (X-1)(X+1)(X^4+X^3+X^2+X+1)(X^4-X^3+X^2-X+1). \tag{4.1$'$}$$

The pairwise partition of the irreducible factors such that in every pair one polynomial is the alternating polynomial of the other one are:

$$\{X-1,\ X+1\},\ \ \{X^4 - X^3 + X^2 - X + 1,\ X^4 + X^3 + X^2 + X + 1\}. \tag{4.2$'$}$$

Select one from each pair as follows:

$$X - 1, \quad X^4 - X^3 + X^2 - X + 1. \tag{4.3$'$}$$

The check polynomial of C is:

$$h(X) = (X-1)(X^4-X^3+X^2-X+1) = X^5-2X^4+2X^3-2X^2+2X-1. \tag{4.4$'$}$$

and $\widetilde{h}(X) = X^5 + 2X^4 + 2X^3 + 2X^2 + 2X + 1$, the check polynomial of $C^\perp$ is:

$$\widetilde{h}^*(X) = X^5 + 2X^4 + 2X^3 + 2X^2 + 2X + 1. \tag{4.5$'$}$$

4.3 A class of iso-self-dual MDS cyclic codes

From Theorem 4.5, we can further construct a class of iso-self-dual MDS cyclic codes. Note that $q \equiv 1 \pmod 4$ if and only if $\nu_2(q+1) = 1$.

Theorem 4.8. *Assume that $q \equiv 1 \pmod 4$, and $n = q + 1$. Let $n' = n/2$ and*

$$P = \Big\{ -\frac{n-2}{4},\ \ \cdots,\ \ -1,\ 0,\ 1,\ \cdots,\ \frac{n-2}{4} \Big\} \ \subseteq\ \mathbb{Z}_n.$$

Then P is a μ_q-invariant subset of $\mathbb{Z}_n$ and

(i) $C_P \subseteq R_n$ is an iso-self-dual cyclic code, and $\varphi_{-1,n'}(C_P) = C_P^\perp$;

(ii) C_P is an alternant code over $\mathbb{F}_q$ of length n by restricting a generalized Reed-Solomon code over $\mathbb{F}_{q^2}$ of length n to the subfield $\mathbb{F}_q$, in particular, C_P is a $[q+1,\ \frac{q+1}{2},\ \frac{q+3}{2}]$ MDS code.

Proof. Note that $|P| = \frac{n}{2} = n'$ is odd,

$$\bar{P} = \mathbb{Z}_n \backslash P = \Big\{ \frac{n+2}{4},\ \frac{n+2}{4} + 1,\ \cdots,\ \frac{3n-2}{4} \Big\},$$

and $q \equiv -1 \pmod{n}$. Then for any $i \in P$, i.e., $-\frac{n-2}{4} \leq i \leq \frac{n-2}{4}$, we have $qi \equiv -i \pmod{n}$, hence

$$\frac{n-2}{4} \geq -i \geq -\frac{n-2}{4};$$

i.e., $qi \in P$. We get that P is μ_q-invariant.

(i). Let $i \in P$, i.e., $-\frac{n-2}{4} \leq i \leq \frac{n-2}{4}$. Note that $\tau_{n'}(i) = i + \frac{n}{2} \pmod{n}$, and $-\frac{n-2}{4} \equiv n - \frac{n-2}{4} = \frac{3n+2}{4} \pmod{n}$. Then

$$\tau_{n'}(i) = i + \frac{n}{2} \geq \frac{n}{2} - \frac{n-2}{4} \geq \frac{n+2}{4} > \frac{n-2}{4};$$

and

$$\tau_{n'}(i) = i + \frac{n}{2} \leq \frac{n}{2} + \frac{n-2}{4} = \frac{3n-2}{4} < \frac{3n+2}{4}.$$

We obtain that $\tau_{n'}(i) \notin P$. So $P \bigcap \tau_{n'}(P) = \emptyset$; and $\mathbb{Z}_n = P \bigcup \tau_{n'}(P)$ is a partition. By Theorem 2.12, (i) holds.

(ii). Let θ be a primitive n-th root of unity. Since $n = q + 1$ divides $q^2 - 1$, we see that $\theta \in \mathbb{F}_{q^2}$. The subset P is of course μ_{q^2}- invariant, By $\tilde{C}_P$ we denote the cyclic code over $\mathbb{F}_{q^2}$ of length n (i.e., $\tilde{C}_P \subseteq \mathbb{F}_{q^2}[X]/\langle X^n - 1\rangle$) with check polynomial

$$f_P(X) = \prod_{i \in P} (X - \theta^i) \in \mathbb{F}_{q^2}[X],$$

hence $f_{\bar{P}}(X) = \prod_{h \in \bar{P}} (X - \theta^h)$ is a generator polynomial of $\tilde{C}_P$. Of course, $\tilde{C}_P$ is iso-self-dual. Denote $\ell = \frac{n-2}{4}$. We claim that

$$\tilde{C}_p = \left\{ \left(a(1), \theta^\ell a(\theta^{-1}), \cdots, \theta^{(n-1)\ell} a(\theta^{-(n-1)})\right) \,\middle|\, a(X) \in \mathbb{F}_{q^2}[X], \deg a(X) < n' \right\} \tag{4.6}$$

where the right hand side is a generalized Reed-Solomon code over $\mathbb{F}_{q^2}$ of length n.

Note that $\{\theta^h \mid h \in \bar{P}\}$ is the defining set of the cyclic code $\tilde{C}_P$. Any word $\left(a(1), \theta^\ell a(\theta^{-1}), \cdots, \theta^{(n-1)\ell} a(\theta^{-(n-1)})\right)$, for $a(X) \in \mathbb{F}_{q^2}[X]$ with $\deg a(X) < n'$, corresponds to the polynomial

$$C_a(X) = \sum_{i=0}^{n-1} \theta^{\ell i} a(\theta^{-i}) X^i \in \mathbb{F}_{q^2}[X].$$

To proof Eqn (4.6), it is enough to show that, for any $h \in \bar{P}$ and any $a(X) \in \mathbb{F}_{q^2}[X]$ with $\deg a(X) < n'$, the θ^h is a root of $C_a(X)$. Write $a(X) = \sum_{j=0}^{n'-1} a_j X^j$. Then

$$C_a(\theta^h) = \sum_{i=0}^{n-1} \theta^{\ell i} a(\theta^{-i}) \theta^{hi} = \sum_{i=0}^{n-1} \sum_{j=0}^{n'-1} a_j \theta^{-ji} \theta^{\ell i} \theta^{hi};$$

i.e.,

$$C_a(\theta^h) = \sum_{j=0}^{n'-1} a_j \sum_{i=0}^{n-1} \theta^{(h-(j-\ell))i}. \tag{4.7}$$

Recall that $\ell = \frac{n-2}{4}$. For $0 \le j \le n' - 1 = \frac{n-2}{2}$, we get

$$-\frac{n-2}{4} \le j - \ell \le \frac{n-2}{2} - \frac{n-2}{4} = \frac{n-2}{4};$$

in other words, $j - \ell \in P$. However, $h \in \bar{P}$. So $j - \ell \not\equiv h \pmod{n}$, equivalently, $h - (j - \ell) \not\equiv 0 \pmod{n}$. Noting that θ is a primitive n-th root of unity, we see that $\theta^{h-(j-\ell)} \neq 1$ and $\theta^n = 1$. Thus,

$$\sum_{i=0}^{n-1} \theta^{(h-(j-\ell))i} = \frac{\theta^{(h-(j-\ell))n} - 1}{\theta^{h-(j-\ell)} - 1} = 0.$$

Return to Eqn (4.7), we obtain that $C_a(\theta^h) = 0$ for all $h \in \bar{P}$. The conclusion of Eqn (4.6) is proved.

Finally, since the subset P of $\mathbb{Z}_n$ is μ_q-invariant, the polynomial $f_P(X)$ is in fact an $\mathbb{F}_q$-polynomial. The cyclic code C_P over $\mathbb{F}_q$ has $f_P(X)$ as a check polynomial, so C_P consists of the codewords of $\tilde{C}_P$ whose coefficients are all in $\mathbb{F}_q$. We are done. □

The performance of the iso-self-dual MDS cyclic codes in Theorem 4.8 would be nice. These codes have the following properties:

- they have good algebraic structure: they are cyclic and iso-self-dual;
- they have good weight structure: they are MDS and formally self-dual;
- they can be systematically encoded by feed-back shift registers;
- they can be decoded by Berlekamp-Welch decoding algorithm.

The following is an instance of the class of codes constructed in Theorem 4.8.

Example 4.9. Let $q = 5$, $n = q + 1 = 6$. Then $q = 5 \equiv 1 \pmod 4$, $\nu_2(n) = 1$ and $n' = 6/2 = 3$. By Theorem 4.8, we take $t = n' = 3$, and

$$P = \{0, 1, -1 = 5\}, \quad \text{hence} \quad \bar{P} = \mathbb{Z}_{10} \backslash P = \{2, 3, 4\}$$

and $\tau_t(P) = \bar{P}$. Take θ to be a primitive 6-th root of unity such that $\theta^3 = -1$. Then C_P is an iso-self-dual MDS cyclic code over $\mathbb{F}_5$ of length 6 with the check polynomial $f_P(X)$ and the generator polynomial $f_{\bar{P}}(X)$ as follows (note that $\theta^3 = -1$):

$$\begin{aligned} f_P(X) &= (X - \theta^0)(X - \theta)(X - \theta^5) = (X - 1)(X^2 - X + 1) \\ &= X^3 - 2X^2 + 2X - 1, \\ f_{\bar{P}}(X) &= (X - \theta^3)(X - \theta^2)(X - \theta^4) = (X + 1)(X^2 + X + 1) \\ &= X^3 + 2X^2 + 2X + 1. \end{aligned}$$

And $C_P^\perp = \varphi_{-1,t}(C_P)$ has a check polynomial $f_{-\bar{P}}(X) = X^3 + 2X^2 + 2X + 1$. By Theorem 4.8(ii) and Eqn (4.6), C_P is the alternant code by restricting the following generalized Reed-Solomon code over $\mathbb{F}_{25}$ of length 6:

$$\big\{\big(a(1), \theta a(\theta^{-1}), \cdots, \theta^5 a(\theta^{-5})\big) \;\big|\; a(X) \in \mathbb{F}_{25}[X],\ \deg a(X) < 3\big\}.$$

Acknowledgements

The research of the authors is supported by NSFC with grant number 11271005.

References

[1] J. L. Alperin, R. B. Bell, Groups and Representations, GTM 162, Springer-Verlag, New York, 1997.

[2] S. A. Aly, A. Klappenecker, P. K. Sarvepalli, Duadic group algebra codes, In Proc. Int. Symp. Inf. Theory, Adelaide, Australia, (2007), 2096-2100.

[3] T. Blackford, Negacyclic duadic codes, Finite Fields Appl., **14**(2008), 930-943.

[4] T. Blackford, Isodual constacyclic codes, Finite Fields Appl., **24**(2013), 29-44.

[5] R. A. Brualdi, V. Pless, Polyadic codes, Discr. Appl. Math., **25**(1989), 3-17.

[6] B. Chen and H. Q. Dinh, A note on isodual constacyclic codes, Finite Fields Appl., vol. 29, pp. 243-246, Sep. 2014.

[7] B. Chen, H. Q. Dinh, Y. Fan, S. Ling, Polyadic constacyclic codes, IEEE Trans. Inform. Theory **61**(2015), no.9, 4895-4904.

[8] B. Chen, Y. Fan, L. Lin, H. Liu, Constacyclic codes over finite fields, Finite Fields Appl., **18**(2012), 1217-1231.

[9] C. Ding, K.Y. Lam, C. Xing, Enumeration and construction of all duadic codes of length p^m, Fund. Inform. **38**(1999), 149-161.

[10] H. Q. Dinh, Lopez-Permouth, Cyclic and negacyclic codes over finite chain rings, IEEE Trans. Inform. Theory **50**(2004), no.8, 1728-1744.

[11] C. Ding, V. Pless, Cyclotomy and duadic codes of prime lengths, IEEE Trans. Inform. Theory, **45**(1999), 453-466.

[12] Y. Fan, G. Zhang, On the existence of self-dual permutation codes of finite groups, Des. Codes Cryptogr., **62**(2012), 19-29.

[13] Y. Fan, L. Zhang, Iso-orthogonality and Type-II duadic constacyclic codes, Finite Fields Appl. **41** (2016) 1-23.

[14] Y. Fan, L. Zhang, Galois self-dual constacyclic codes, Des. Codes Cryptogr., DOI 10.1007/s10623-016-0282-8, online: 29 Sept. 2016.

[15] S. Han, J-L. Kim, Computational results of duadic double circulant codes, J. Appl. Math. Comput., **40**(2012), 33-43.

[16] W.C. Huffman, V. Pless, Fundamentals of Error-Correcting Codes, Cambridge University Press, Cambridge, 2003.

[17] S. Jitman, S. Ling, P. Solé, Hermitian self-dual Abelian codes, IEEE Trans. Inform. Theory, 60(2014),1496-1507.

[18] J. S. Leon, J. M. Masley, V. Pless, Duadic codes, IEEE Trans. Inform. Theory, **30**(1984), 709-714.

[19] S. Ling, C. Xing, Polyadic codes revisited, IEEE Trans. Inform. Theory, 50(2004),200-207.

[20] C. Martinez-Pérez, W. Willems, Self-dual codes and modules for finite groups in characteristic two, IEEE Trans. Inform. Theory 50(2004), no. 8, 1798-1803.

[21] V. Pless, Duadic codes revisited, Congressus Numeratium, **59**(1987), 225-233.

[22] J. J. Rushanan, Duadic codes and difference sets, J. Combin. Theory Set. A, **57**(1991), 254-61.

[23] A. Sharma, G.K. Bakshi, V.C. Dumir, M. Raka, Cyclotomic numbers and primitive idempotents in the ring $GF(q)[X]/(X^{p^n}-1)$, Finite Fields Appl. 10 (4) (2004) 653-673.

[24] M. H. M. Smid, Duadic codes, IEEE Trans. Inform. Theory, **33**(1987), 432-433.

[25] H. N. Ward, L. Zhu, Existence of abelian group codes partitions, J. Combin. Theory Ser. A, **67**(1994), 276-281.

[26] W. Willems, A note on self-dual group codes, IEEE Trans. Inform. Theory 48(2002), 3107-3109.